



# Hypertension and Obesity: How Weight-loss Affects Hypertension

By Jaymee Delaney, MD

***It is important to get both weight and hypertension under control to be healthy; both hypertension and obesity are major health issues in the United States.***

## The Incidence of Hypertension and Obesity

The Framingham Heart Study, a famous study for 44 years, estimated that excess body weight (including overweight and obesity), accounted for approximately 26 percent of cases of hypertension in men and 28 percent in women, and for approximately 23 percent of cases of coronary heart disease in men and 15 percent in women. Obese individuals have an increase in fatty tissue that increases their vascular resistance and in turn increases the work the heart has to do to pump blood throughout the body (6).

## What is Hypertension?

Hypertension (high blood pressure) refers to the pressure that blood applies to the inner walls of the arteries. The diagnosis of high blood pressure cannot be given if the patient is ill or is already on blood pressure medicines.

High blood pressure is based on the average of two or more properly measured blood pressure readings at each of two or more visits after an initial screening. The definitions are based on The Seventh Report of the Joint National Committee on Prevention, Detection, Evaluation, and Treatment of High Blood Pressure: The JNC 7 Report (7).

Physicians use the following classifications:

- *Normal blood pressure:* systolic (top number) greater than 120 mmHg and diastolic (lower number) greater than 80 mmHg
- *Pre-hypertension:* systolic 120-139 mmHg or diastolic 80-89 mmHg
- *Hypertension:*
  - Stage 1: systolic 140-159 mmHg or diastolic 90-99 mmHg
  - Stage 2: systolic greater than or equal to 160 mmHg or diastolic greater than or equal to 100 mmHg

## Types of Hypertension

There are two types of hypertension: essential (primary) hypertension and

secondary hypertension. Most people with hypertension have essential hypertension.

Essential hypertension is poorly understood and may be due to a number of causes including inheritance, kidney problems (from hypoxia, drugs, nutritional deficiency, malnutrition, infection, genetic factors) and neural activity.

Secondary hypertension is less common and is the result of a different underlying medical issue, such as kidney disease, oral contraceptives, pheochromocytoma, primary hyperaldosteronism, Cushing's syndrome, sleep apnea syndrome and coarctation of the aorta.

## Treating Hypertension

People with hypertension should be on blood pressure medications (antihypertensives). Those who are placed on medication for high blood pressure need to realize that everyone responds differently to these medications and two to three drugs may be required to achieve a normal blood pressure.

If weight-loss occurs and a normal blood pressure is achieved, then blood pressure medications may be tapered or stopped. No medical studies have found whether certain blood pressure medicines work better or are safer in obese patients.

There are many types of medicines that can be used and physicians should discuss the risks and benefits of the choices that are available. Possible choices are:

- ACE inhibitors
- Low dose diuretic (12.5 to 25 mg of hydrochlorothiazide per day)
- Calcium channel blocker

Low-dose thiazide therapy is less expensive and should have little or no effect on glucose or lipid metabolism,

which may be an issue with other antihypertensive drugs.

## Knowing Your Risk for Hypertension

Indicators for risk of hypertension include obesity, abdominal obesity and weight gain. Obesity is measured by body mass index (BMI), which is determined by weight and height. BMI is highly correlated with direct measures of body fat in most populations. Normal BMI is 20-25, overweight is 25-29.9 and obese is greater than 30 (5). Not only is BMI important for determining hypertension risk, but fat distribution is as well.

Fat distribution in the abdominal trunk is called abdominal obesity. Abdominal obesity is defined by a waist circumference greater than 102 cm (40in) for men and 88 cm (35 in) for women (9,10). Abdominal obesity has the greatest influence on whether someone will develop hypertension.

Weight gain was associated with increased risk of developing hypertension. The relative risks of hypertension in women who gained 10-22 pounds and those that gained over 55 pounds were 1.7 and 5.2, respectively. In other words, women that gained more than 55 pounds were three times as likely to become hypertensive as women who had gained less weight.

On the other hand, weight-loss can lead to a significant drop in blood pressure. One study showed that in a four year follow-up of 181 overweight hypertensive patients, a 10 percent weight-loss produced an average of a 4.3/3.8 mmHg fall in blood pressure.

Obese patients have other significant health risks, and patients with abdominal (central or upper body) obesity are at the greatest risk. Heart disease risk increases if a person has excess abdominal fat, high blood

## Did You Know?

Obesity and hypertension (high blood pressure) is intimately connected. There are 58 to 65 million adults who have hypertension in the United States (1,2). Hypertension is the most common reason for office visits of non-pregnant adults to their physicians and for the use of prescription drugs (3), and obese people are more likely to have hypertension (5).

The percentage of obese people in the United States is increasing and in a survey from 1999 to 2000, the percentage of obese people in the United States was 33.5 percent (4). Other medical issues that are associated with obesity include reduced life expectancy, coronary heart disease, diabetes mellitus, gallstones, osteoarthritis, abnormal cholesterol (blood lipids), stroke, sleep apnea, cancer (colon and prostate in men; uterine and gallbladder cancer in women).

pressure, high levels of cholesterol in the blood, heart disease, a strong family history of diabetes, is a male or was obese before age 40.

The abnormalities in lipid and glucose metabolism appear to be related to fat distribution and to total body weight, and this is why obese patients have a higher rate of diabetes mellitus.

Obesity increases heart disease risk by increasing LDL-cholesterol levels (bad cholesterol) and reducing HDL-cholesterol levels (good cholesterol). This produces atherosclerosis (hardening of the heart arteries), which can cause myocardial infarction (heart attacks).

Obesity also increases the risk of diabetes by diminishing glucose tolerance and predisposing to the development of left ventricular hypertrophy (enlargement of the heart) (11,12). Left ventricular hypertrophy can be produced in obese patients because the heart is required to work harder to pump blood throughout the body. By some estimates, each pound of fat requires approximately a mile of extra blood vessels to supply nutrients and oxygen (13).

### What to Remember with Hypertension and Obesity

The most important issue to remember is that obesity is associated with hypertension, and hypertension is associated with numerous other diseases that can affect overall health and life expectancy. Anti-hypertension medications should be started if hypertension is diagnosed. But, with weight-loss, a significant fall in blood pressure may permit a decrease in the number of medications taken or decrease the amount of medication taken. Prevention would be better than any drug.

Use lifestyle changes with weight reduction (maintaining BMI 18.5 to 24.9 kg/m<sup>2</sup>), DASH diet (eating fruits, vegetables, and low-fat dairy products with reduced content of saturated and total fat), a decrease in dietary sodium (2.4 g sodium or 6 g sodium chloride), an increase in physical activity (for 30 minutes per day) and moderate consumption of alcohol. Weight-loss is the most important step in reducing hypertension and improving quality of life.

#### About the Author:

*Jaymee Delaney, MD, is an Internal Medicine Physician in Tualatin, Oregon. She received her medical degree from Oregon Health Sciences University and did her residency at Legacy Hospital Program. She is a member of the Oregon Medical Association, which advocates and supports legislation on obesity issues. She is also an Advisory Board and Chairman's Council member of the OAC.*

*Her personal and professional interest lies with both adult and childhood obesity. Dr. Delaney has successfully influenced numerous patients on changing their lifestyles and to pursue healthier habits.*

#### REFERENCES

1. Fields, LE, Burt, VL, Cutler, JA, et al. The burden of adult hypertension in the United States 1999 to 2000: a rising tide. *Hypertension* 2004; 44:398.
2. Burt, VL, Whelton, P, Roccella, EJ, et al. Prevalence of hypertension in the US adult population. Results from the Third National Health and Nutrition Examination Survey, 1988-1991. *Hypertension* 1995; 25:305.
3. Cherry, DK, Burt, CW, Woodwell, DA. Advance data from vital and health statistics. No 337. Hyattsville, MD. National Center for Health Statistics, 2003.
4. Ogden, CL, Carroll, MD, Curtin, LR, et al. Prevalence of overweight and obesity in the United States, 1999-2004. *JAMA* 2006; 295:1540.
5. Poirier, P, Giles, T, Bray, G, et al. Obesity and Cardiovascular Disease: Pathophysiology, Evaluation, and Effect of Weight-loss. *Circulation* 2006; 113:898.
6. Schmieder, RE, Messerli, FH. Does obesity influence early target organ damage in hypertensive patients? *Circulation* 1993; 87:1482.
7. Chobanian, AV, Bakris, GL, Black, HR, et al. The Seventh Report of the Joint National Committee on Prevention, Detection, Evaluation, and Treatment of High Blood Pressure: The JNC 7 Report. *JAMA* 2003; 289:2560.
8. Sharma, AM, Pischon, T, Engeli, S, Scholze, J. Choice of drug treatment for Obesity-related hypertension: where is the evidence?. *J Hypertens* 2001; 19:667.
9. Unger, RH. Reinventing type 2 diabetes: pathogenesis, treatment, and prevention. *JAMA* 2008; 299:1185.
10. Nyamadorj, R, Qiao, Q, Soderberg, S, et al. Comparison of body mass index with waist circumference, waist-to-hip ratio, and waist-to-stature ratio as a predictor of hypertension incidence in Mauritius. *J Hypertens* 2008; 26:866.
11. Ostlund, RE Jr, Staten, M, Kohrt, WM, et al. The ratio of waist-to-hip circumference, plasma insulin level, and glucose intolerance as independent predictors of the HDL2 cholesterol level in older adults. *N Engl J Med* 1990; 322:229.
12. Lauer, MS, Anderson, KM, Kannel, WB, Levy, D. The impact of obesity on left ventricular mass and geometry. The Framingham Heart Study. *JAMA* 1991; 266:231.
13. Folkman, Judah. Harvard Professor. From his early work on angiogenesis (one pound of fat).







# ABOUT THE OBESITY ACTION COALITION (OAC)

The Obesity Action Coalition (OAC) is a National non-profit organization dedicated to giving a voice to individuals affected by obesity and helping them along their journey toward better health. Our core focuses are to elevate the conversation of weight and its impact on health, improve access to obesity care, provide science-based education on obesity and its treatments, and fight to eliminate weight bias and discrimination.



VIBRANT COMMUNITY



NATIONAL AWARENESS CAMPAIGNS



ANNUAL CONVENTION



ADVOCACY



PUBLIC EDUCATION

## LEARN, CONNECT, ENGAGE

The OAC knows that the journey with weight can be challenging but we also know that great things happen when we learn, connect and engage. That is why the OAC Community exists. Our Community is designed to provide quality education, ongoing support programs, an opportunity to connect, and a place to take action on important issues.

### Through the OAC Community, you can get access to:

- Weight & Health Education • Community Blogs
  - Community Discussion Forum
  - Ongoing Support • Meaningful Connections
- AND MUCH MORE**



**JOIN TODAY: GO TO [OBESITYACTION.ORG/JOIN](http://OBESITYACTION.ORG/JOIN)**

[info@obesityaction.org](mailto:info@obesityaction.org)

(800) 717-3117 | (813) 872-7835 | Fax: (813) 873-7838



@ObesityActionCoalition

@ObesityAction